

QUADRATIC TRANSFORMATIONS AND GUILLERA'S FORMULAE FOR $1/\pi^2$

WADIM ZUDILIN

ABSTRACT. We prove two new series of Ramanujan type for $1/\pi^2$.

In a series of works [3]–[5], J. Guillera discovered several Ramanujan-type formulae for $1/\pi^2$, some of which he proved. For instance, the following two identities are proven in [3], [5]:

$$\sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^5}{n!^5} (20n^2 + 8n + 1) \left(-\frac{1}{2^2}\right)^n = \frac{8}{\pi^2}; \quad (1)$$

$$\sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^5}{n!^5} (820n^2 + 180n + 13) \left(-\frac{1}{2^{10}}\right)^n = \frac{128}{\pi^2}, \quad (2)$$

where $(a)_n = \Gamma(a+n)/\Gamma(a) = \prod_{k=0}^{n-1} (a+k)$ denotes Pochhammer's symbol. It happens that, in all Guillera's formulae expressions, the left-hand sides are given by certain hypergeometric ${}_5F_4$ -series, with the sole exception [6]

$$\sum_{n=0}^{\infty} A_n \frac{36n^2 + 12n + 1}{2^{10n}} = \frac{32}{\pi^2}, \quad \text{where } A_n = \binom{2n}{n}^2 \sum_{k=0}^n \binom{2k}{k}^2 \binom{2n-2k}{n-k}^2 \quad (3)$$

(this formula remains unproven). Recall that a generalized hypergeometric series [7] is defined for $z \in \mathbb{C}$, $|z| < 1$, by

$${}_qF_q \left(\begin{matrix} a_0, a_1, \dots, a_q \\ b_1, \dots, b_q \end{matrix} \middle| z \right) = \sum_{n=0}^{\infty} \frac{(a_0)_n (a_1)_n \cdots (a_q)_n}{n! (b_1)_n \cdots (b_q)_n} z^n.$$

Formula (3) looks a natural extension of Sato's formula for $1/\pi$ involving Apéry's numbers (see [9] and [2]) as well as of several other examples like

$$\sum_{n=0}^{\infty} B_n \frac{4n+1}{36^n} = \frac{18}{\pi\sqrt{15}}, \quad \text{where } B_n = \sum_{k=0}^n \binom{n}{k}^4,$$

proven by Y. Yang [10].

The aim of this note is to derive two new identities of type (3) from (1), (2) using hypergeometric technique.

1. QUADRATIC TRANSFORMATIONS

Recall quadratic transformations for ${}_2F_1$ -series,

$${}_2F_1 \left(\begin{matrix} a, b \\ 1+a-b \end{matrix} \middle| z \right) = (1-z)^{-a} \cdot {}_2F_1 \left(\begin{matrix} \frac{1}{2}a, \frac{1}{2} + \frac{1}{2}a - b \\ 1+a-b \end{matrix} \middle| \frac{-4z}{(1-z)^2} \right)$$

due to Gauss, and for ${}_3F_2$ -series,

$${}_3F_2 \left(\begin{matrix} a, b, c \\ 1+a-b, 1+a-c \end{matrix} \middle| z \right) = (1-z)^{-a} \cdot {}_3F_2 \left(\begin{matrix} \frac{1}{2}a, \frac{1}{2} + \frac{1}{2}a, 1+a-b-c \\ 1+a-b, 1+a-c \end{matrix} \middle| \frac{-4z}{(1-z)^2} \right) \quad (4)$$

Date: April 2005.

The work is supported by a fellowship of the Max-Planck Institute for Mathematics (Bonn).

due to Whipple. Their higher-order analogues necessarily involve multiple hypergeometric series in the right-hand side (cf., e.g., [1], Proposition 6, and [11], Theorem 5). Here we indicate the following result.

Theorem 1. *The following quadratic transformation is valid:*

$$\begin{aligned} & {}_5F_4\left(\begin{matrix} a, b, c, d, e \\ 1+a-b, 1+a-c, 1+a-d, 1+a-e \end{matrix} \middle| z\right) \\ &= (1-z)^{-a} \sum_{n=0}^{\infty} \frac{(\frac{1}{2}a)_n (\frac{1}{2} + \frac{1}{2}a)_n}{(1+a-b)_n (1+a-c)_n} \left(\frac{-4z}{(1-z)^2}\right)^n \\ &\quad \times \sum_{\nu=0}^n \frac{(b)_{\nu} (c)_{\nu} (1+a-d-e)_{\nu}}{\nu! (1+a-d)_{\nu} (1+a-e)_{\nu}} \frac{(1+a-b-c)_{n-\nu}}{(n-\nu)!}, \end{aligned} \quad (5)$$

whenever both series converge.

Remark. Theorem 1 may be stated in the form of Orr-type theorem (cf. [7], Section 2.5): If

$$(1-z)^{b+c-a-1} \cdot {}_3F_2\left(\begin{matrix} b, c, 1+a-d-e \\ 1+a-d, 1+a-e \end{matrix} \middle| z\right) = \sum_{n=0}^{\infty} f_n z^n, \quad (6)$$

then

$$\begin{aligned} & {}_5F_4\left(\begin{matrix} a, b, c, d, e \\ 1+a-b, 1+a-c, 1+a-d, 1+a-e \end{matrix} \middle| z\right) \\ &= (1-z)^{-a} \sum_{n=0}^{\infty} f_n \frac{(\frac{1}{2}a)_n (\frac{1}{2} + \frac{1}{2}a)_n}{(1+a-b)_n (1+a-c)_n} \left(\frac{-4z}{(1-z)^2}\right)^n. \end{aligned}$$

It follows from (6) that $|f_n|^{1/n} \rightarrow 1$ as $n \rightarrow \infty$, hence the condition $|4z/(1-z)^2| < 1$ ensures convergence of the double series in (5).

Proof. It is sufficient to prove the identity in a neighbourhood of the origin. Using the Pfaff–Saalschütz theorem [7], formula (2.3.1.3),

$${}_3F_2\left(\begin{matrix} -n, a+n, 1+a-d-e \\ 1+a-d, 1+a-e \end{matrix} \middle| 1\right) = \frac{(-d-n+1)_n (e)_n}{(1+a-d)_n (e-a-n)_n} = \frac{(d)_n (e)_n}{(1+a-d)_n (1+a-e)_n},$$

write

$$\begin{aligned} & {}_5F_4\left(\begin{matrix} a, b, c, d, e \\ 1+a-b, 1+a-c, 1+a-d, 1+a-e \end{matrix} \middle| z\right) \\ &= \sum_{n=0}^{\infty} \frac{(a)_n (b)_n (c)_n}{n! (1+a-b)_n (1+a-c)_n} z^n \cdot {}_3F_2\left(\begin{matrix} -n, a+n, 1+a-d-e \\ 1+a-d, 1+a-e \end{matrix} \middle| 1\right) \\ &= \sum_{n=0}^{\infty} \frac{(a)_n (b)_n (c)_n}{n! (1+a-b)_n (1+a-c)_n} z^n \sum_{\nu=0}^n \frac{(-n)_{\nu} (a+n)_{\nu} (1+a-d-e)_{\nu}}{\nu! (1+a-d)_{\nu} (1+a-e)_{\nu}} \\ &= \sum_{\nu=0}^{\infty} \frac{(1+a-d-e)_{\nu} (-1)^{\nu}}{\nu! (1+a-d)_{\nu} (1+a-e)_{\nu}} \sum_{n=\nu}^{\infty} \frac{(a)_{n+\nu} (b)_n (c)_n}{(n-\nu)! (1+a-b)_n (1+a-c)_n} z^n \\ &= \sum_{\nu=0}^{\infty} \frac{(1+a-d-e)_{\nu} (-1)^{\nu}}{\nu! (1+a-d)_{\nu} (1+a-e)_{\nu}} \sum_{m=\nu}^{\infty} \frac{(a)_{m+2\nu} (b)_{m+\nu} (c)_{m+\nu}}{m! (1+a-b)_{m+\nu} (1+a-c)_{m+\nu}} z^{m+\nu} \end{aligned}$$

$$\begin{aligned}
 &= \sum_{\nu=0}^{\infty} \frac{(a)_{2\nu}(b)_{\nu}(c)_{\nu}(1+a-d-e)_{\nu}}{\nu!(1+a-b)_{\nu}(1+a-c)_{\nu}(1+a-d)_{\nu}(1+a-e)_{\nu}} (-z)^{\nu} \\
 &\quad \times {}_3F_2\left(\begin{matrix} a+2\nu, b+\nu, c+\nu \\ 1+a-b+\nu, 1+a-c+\nu \end{matrix} \middle| z\right).
 \end{aligned}$$

Applying the quadratic transformation (4) to the latter ${}_3F_2$ -series, we deduce

$$\begin{aligned}
 &{}_5F_4\left(\begin{matrix} a, b, c, d, e \\ 1+a-b, 1+a-c, 1+a-d, 1+a-e \end{matrix} \middle| z\right) \\
 &= \sum_{\nu=0}^{\infty} \frac{(a)_{2\nu}(b)_{\nu}(c)_{\nu}(1+a-d-e)_{\nu}}{\nu!(1+a-b)_{\nu}(1+a-c)_{\nu}(1+a-d)_{\nu}(1+a-e)_{\nu}} (-z)^{\nu} \\
 &\quad \times (1-z)^{-(a+2\nu)} \cdot {}_3F_2\left(\begin{matrix} \frac{1}{2}a+\nu, \frac{1}{2}+\frac{1}{2}a+\nu, 1+a-b-c \\ 1+a-b+\nu, 1+a-c+\nu \end{matrix} \middle| \frac{-4z}{(1-z)^2}\right) \\
 &= (1-z)^{-a} \sum_{\nu=0}^{\infty} \frac{(\frac{1}{2}a)_{\nu}(\frac{1}{2}+\frac{1}{2}a)_{\nu}(b)_{\nu}(c)_{\nu}(1+a-d-e)_{\nu}}{\nu!(1+a-b)_{\nu}(1+a-c)_{\nu}(1+a-d)_{\nu}(1+a-e)_{\nu}} \left(\frac{-4z}{(1-z)^2}\right)^{\nu} \\
 &\quad \times \sum_{m=0}^{\infty} \frac{(\frac{1}{2}a+\nu)_m(\frac{1}{2}+\frac{1}{2}a+\nu)_m(1+a-b-c)_m}{m!(1+a-b+\nu)_m(1+a-c+\nu)_m} \left(\frac{-4z}{(1-z)^2}\right)^m \\
 &= (1-z)^{-a} \sum_{\nu=0}^{\infty} \frac{(b)_{\nu}(c)_{\nu}(1+a-d-e)_{\nu}}{\nu!(1+a-d)_{\nu}(1+a-e)_{\nu}} \left(\frac{-4z}{(1-z)^2}\right)^{\nu} \\
 &\quad \times \sum_{n=\nu}^{\infty} \frac{(\frac{1}{2}a)_n(\frac{1}{2}+\frac{1}{2}a)_n(1+a-b-c)_{n-\nu}}{(n-\nu)!(1+a-b)_n(1+a-c)_n} \left(\frac{-4z}{(1-z)^2}\right)^{n-\nu} \\
 &= (1-z)^{-a} \sum_{n=0}^{\infty} \frac{(\frac{1}{2}a)_n(\frac{1}{2}+\frac{1}{2}a)_n}{(1+a-b)_n(1+a-c)_n} \left(\frac{-4z}{(1-z)^2}\right)^n \\
 &\quad \times \sum_{\nu=0}^n \frac{(b)_{\nu}(c)_{\nu}(1+a-d-e)_{\nu}}{\nu!(1+a-d)_{\nu}(1+a-e)_{\nu}} \frac{(1+a-b-c)_{n-\nu}}{(n-\nu)!},
 \end{aligned}$$

which is the required formula (5). □

Plugging in $a = b = c = d = e = \frac{1}{2}$ we obtain

$$\sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^5}{n!^5} z^n = \frac{1}{(1-z)^{1/2}} \sum_{n=0}^{\infty} \frac{(\frac{1}{4})_n(\frac{3}{4})_n}{n!^2} \left(\frac{-4z}{(1-z)^2}\right)^n \sum_{\nu=0}^n \frac{(\frac{1}{2})_{\nu}^3}{\nu!^3} \frac{(\frac{1}{2})_{n-\nu}}{(n-\nu)!}.$$

Note the equality

$$u_n = \sum_{\nu=0}^n \frac{(\frac{1}{2})_{\nu}^3}{\nu!^3} \frac{(\frac{1}{2})_{n-\nu}}{(n-\nu)!} = \sum_{\nu=0}^n \left(\frac{(\frac{1}{4})_{\nu}(\frac{3}{4})_{n-\nu}}{\nu!(n-\nu)!} \right)^2, \quad (7)$$

followed from [7], formula (2.5.18). Another way to prove (7) is based on the recurrence relation

$$8(n+1)^3 u_{n+1} - (2n+1)(8n^2 + 8n + 5)u_n + 8n^3 u_{n-1} = 0 \quad (8)$$

satisfied by both expressions in (7). (Proof of (8) uses the algorithm of creative telescoping [8], Chapter 6.) We summarize saying above in

Theorem 2. Suppose $|z| < 1$ and $|4z/(1-z)^2| < 1$. The following identity holds:

$$\sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n^5}{n!^5} z^n = \frac{1}{(1-z)^{1/2}} \sum_{n=0}^{\infty} u_n \frac{(\frac{1}{4})_n(\frac{3}{4})_n}{n!^2} \left(\frac{-4z}{(1-z)^2}\right)^n, \quad (9)$$

where u_n are given in (7).

Remark. It is worth mentioning the following curious hypergeometric identity:

$$\sum_{n=0}^{\infty} u_n \frac{(\frac{1}{3})_n (\frac{2}{3})_n}{n!^2} z^n = {}_3F_2 \left(\frac{1}{6}, \frac{1}{2}, \frac{5}{6} \middle| z \right)^2 = {}_2F_1 \left(\frac{1}{12}, \frac{5}{12} \middle| z \right)^4, \quad (10)$$

since the double hypergeometric series in the left-hand side is very close to the one used in (9). Formula (10) was born in our correspondence with G. Almkvist and Y. Yang.

2. NEW FORMULAE FOR $1/\pi^2$

For the differential operator $\theta = z \frac{d}{dz}$, we have

$$\theta(1-z)^{-1/2} = \frac{z}{2(1-z)} \cdot (1-z)^{-1/2} \quad \text{and} \quad \theta \left(\frac{-4z}{(1-z)^2} \right) = \frac{1+z}{1-z} \cdot \left(\frac{-4z}{(1-z)^2} \right).$$

Therefore,

$$\begin{aligned} & \theta \left((1-z)^{-1/2} \sum_{n=0}^{\infty} C_n \left(\frac{-4z}{(1-z)^2} \right)^n \right) \\ &= (1-z)^{-1/2} \sum_{n=0}^{\infty} C_n \cdot \left(n \frac{1+z}{1-z} + \frac{z}{2(1-z)} \right) \cdot \left(\frac{-4z}{(1-z)^2} \right)^n, \\ & \theta^2 \left((1-z)^{-1/2} \sum_{n=0}^{\infty} C_n \left(\frac{-4z}{(1-z)^2} \right)^n \right) \\ &= (1-z)^{-1/2} \sum_{n=0}^{\infty} C_n \cdot \left(n^2 \frac{(1+z)^2}{(1-z)^2} + n \frac{z(3+z)}{(1-z)^2} + \frac{z(2+z)}{4(1-z)^2} \right) \cdot \left(\frac{-4z}{(1-z)^2} \right)^n. \end{aligned}$$

We now apply the results to the functions

$$20\theta^2 f(z) + 8\theta f(z) + f(z) \quad \text{and} \quad 820\theta^2 f(z) + 180\theta f(z) + 13f(z),$$

where $f(z)$ is given in (9), and substitute $z = -1/2^2$ and $z = -1/2^{10}$, respectively. Using then evaluations (1), (2) and formulae

$$\frac{(\frac{1}{4})_n (\frac{3}{4})_n}{n!^2} = 2^{-6n} \frac{(4n)!}{n!^2 (2n)!}, \quad u_n = \sum_{k=0}^n \frac{(\frac{1}{2})_k^3}{k!^3} \frac{(\frac{1}{2})_{n-k}}{(n-k)!} = 2^{-6n} \sum_{k=0}^n \binom{2k}{k}^3 \binom{2n-2k}{n-k} 2^{4(n-k)},$$

we arrive at

Theorem 3. *The following identities are valid:*

$$\begin{aligned} & \sum_{n=0}^{\infty} U_n \frac{(4n)!}{n!^2 (2n)!} \frac{18n^2 - 10n - 3}{(2^8 5^2)^n} = \frac{10\sqrt{5}}{\pi^2}, \\ & \sum_{n=0}^{\infty} U_n \frac{(4n)!}{n!^2 (2n)!} \frac{1046529n^2 + 227104n + 16032}{(5^4 41^2)^n} = \frac{5^4 41 \sqrt{41}}{\pi^2}, \end{aligned}$$

where the sequence of integers

$$U_n = \sum_{k=0}^n \binom{2k}{k}^3 \binom{2n-2k}{n-k} 2^{4(n-k)}, \quad n = 0, 1, 2, \dots,$$

satisfies the recurrence relation

$$(n+1)^3 U_{n+1} - 8(2n+1)(8n^2 + 8n + 5)U_n + 4096n^3 U_{n-1} = 0, \quad n = 1, 2, \dots$$

It seems quite likely that the hypergeometric machinery could lead to several other formulae for $1/\pi^2$.

Acknowledgments. I thank Gert Almkvist, Christian Krattenthaler and Yi-Fan Yang for constructive comments. The final part of the work was done during my visit of the Max-Planck Institute for Mathematics in Bonn. I thank the staff of the institute for the wonderful working conditions I experienced there.

REFERENCES

- [1] *G. Almkvist, W. Zudilin*, Differential equations, mirror maps and zeta values // Calabi–Yau Varieties and Mirror Symmetry. The BIRS workshop volume (06–11 December 2003), eds. J. Lewis, S.-T. Yau and N. Yui. Providence, RI: Amer. Math. Soc. & International Press, 2006 (to appear); `math.NT/0402386`.
- [2] *H. H. Chan, S. H. Chan, Z. Liu*, Domb’s numbers and Ramanujan–Sato type series for $1/\pi$ // Adv. Math. 2004. V. 186, no. 2. P. 396–410.
- [3] *J. Guillera*, Some binomial series obtained by the WZ-method // Adv. in Appl. Math. 2002. V. 29, no. 4. P. 599–603.
- [4] *J. Guillera*, About a new kind of Ramanujan-type series // Experiment. Math. 2003. V. 12, no. 4. P. 507–510.
- [5] *J. Guillera*, Generators of some Ramanujan’s formulas // Ramanujan J. 2006. V. 11, no. 1. P. 41–48.
- [6] *J. Guillera*, My formulas for $1/\pi^2$ // Manuscript at <http://personal.auna.com/jguillera/pi-formulas.pdf>. 2005.
- [7] *L. J. Slater*, Generalized hypergeometric functions. Cambridge: Cambridge University Press, 1966.
- [8] *M. Petkovšek, H. S. Wilf, D. Zeilberger*, $A = B$. Wellesley, MA: A.K. Peters, Ltd., 1996.
- [9] *Y. Yang*, On differential equations satisfied by modular forms // Math. Z. 2004. V. 246, no. 1–2. P. 1–19.
- [10] *Y. Yang*, Personal communication (August 2005).
- [11] *W. Zudilin*, Well-poised hypergeometric transformations of Euler-type multiple integrals // J. London Math. Soc. 2004. V. 70, no. 1. P. 215–230.

DEPARTMENT OF MECHANICS AND MATHEMATICS
 MOSCOW LOMONOSOV STATE UNIVERSITY
 VOROBIOVY GORY, GSP-2
 119992 MOSCOW, RUSSIA
 URL: <http://wain.mi.ras.ru/>
 E-mail address: `wadim@ips.ras.ru`